

BOUNDARY SLOPE DIAMETER AND CROSSING NUMBER OF 2-BRIDGE KNOTS

THOMAS W. MATTMAN, GABRIEL MAYBRUN & KRISTIN ROBINSON

ABSTRACT. We prove that for 2-bridge knots, the diameter, D , of the set of boundary slopes is twice the crossing number, c . This constitutes partial verification of a conjecture that, for all knots in S^3 , $D \leq 2c$.

1. INTRODUCTION

Ichihara [5] told us of a conjecture for knots in S^3 . Let $D(K)$ denote the diameter of the set of boundary slopes of a knot K and $c(K)$ be the crossing number.

Conjecture 1. *For K a knot in S^3 , $D(K) \leq 2c(K)$.*

(To be precise, Ichihara proposed the conjecture only for Montesinos knots and he and Mizushima [6] have recently given a proof of that case.)

Since 0, being the slope of a Seifert surface, is always included in the set of boundary slopes, we have, as an immediate consequence, a conjecture due to Ishikawa and Shimokawa [7]:

Conjecture 2. *Let b be a finite boundary slope for K a knot in S^3 . Then $|b| \leq 2c(K)$.*

For example, it is easy to verify these conjectures for torus knots. For the unknot, $D(K) = 0 = 2c(K)$. For a non-trivial torus knot $K = (p, q)$ we can assume p, q relatively prime with $2 \leq q < p$. The boundary slopes are 0 and pq [9] while the crossing number is $c(K) = pq - p$ [10]. Thus, $D(K) = pq \leq pq + p(q - 2) = 2c(K)$. Moreover, we have equality for the torus 2-bridge knots which are of the form $(p, 2)$ with p odd.

We will show that this equality obtains for all 2-bridge knots:

Theorem 1. *For K a 2-bridge knot, $D(K) = 2c(K)$.*

Corollary 1. *Let b be a boundary slope for a 2-bridge knot K . Then $|b| \leq 2c(K)$.*

This bound is sharp for the $(p, 2)$ torus knots and there are many examples showing that it is also sharp for hyperbolic 2-bridge knots.

Using Conway notation, we can associate a rational number p/q to each 2-bridge knot $K = K(p/q)$. Hatcher and Thurston [4] showed how to calculate the boundary

1991 *Mathematics Subject Classification.* Primary 57M25, 57M27.

Key words and phrases. boundary slopes, crossing number, 2-bridge knots, rational knots.

The second and third authors are undergraduate students who worked under the supervision of the first author during an REU at CSU, Chico in the Summer of 2005. The second author was funded through NSF REU Award 0354174. The third author received support from the MAA's program for Strengthening Underrepresented Minority Mathematics Achievement (SUMMA) with funding from the NSF, NSA, and Moody's. The first author received funding through both programs.

slopes of K from continued fraction representations of p/q . On the other hand, the crossing number is given by summing the terms in a simple continued fraction for p/q (see [3]). Our technique is, starting with a simple continued fraction for p/q , to compute all possible boundary slope continued fractions and identify those which yield the maximum and minimum boundary slopes. We can then verify that the difference between the maximum and minimum boundary slopes is twice the crossing number.

In Section 4 we develop four identities for continued fractions and in Section 5, we use those identities to establish four substitution rules. These substitution rules will allow us to produce all possible boundary slope continued fractions for a given rational number:

Theorem 2. *The boundary slope continued fractions of $K(p/q)$ are among the continued fractions obtained by applying substitutions at non-adjacent positions in the simple continued fraction of p/q .*

The proof of Theorem 2 is presented in Section 6 along with the following corollary.

Corollary 2. *If $\frac{p}{q} = [0, a_0, a_1, \dots, a_n]$ is a simple continued fraction, then $K(p/q)$ has at most F_{n+2} boundary slopes where F_n is the n th Fibonacci number.*

In Section 7 we outline our method for calculating the maximum and minimum boundary slopes and in Section 8 we prove Theorem 1.

Let us now review the basic ideas of Conway notation and continued fractions (Section 2) and boundary slopes for 2-bridge knots (Section 3).

2. CONWAY NOTATION & CONTINUED FRACTIONS

In this section, we give a brief overview of Conway notation, continued fractions, and their relationship.

Rational tangles may be constructed by means of tangle algebra (for example, see Adams [1]). A rational knot is obtained from numerator closure on a rational tangle. The *Conway notation* for such a knot is the same as the Conway notation for the tangle, i.e., a list of integers, $a_0 a_1 \dots a_n$. Note that the set of rational knots coincides with the set of 2-bridge knots and we will use these terms interchangeably.

A *continued fraction* of a rational number $\frac{p}{q}$ is a fraction of the form

$$\frac{p}{q} = c + \frac{1}{b_0 + \frac{1}{b_1 + \frac{1}{\dots + \frac{1}{b_n}}}} = [c, b_0, b_1, \dots, b_n],$$

where $c \in \mathbb{Z}$ and each b_i , for $0 \leq i \leq n$, is a nonzero integer. Note that, since each b_i can be any nonzero integer, the continued fraction for $\frac{p}{q}$ is not unique. We will call c the *integral component* of the continued fraction, and each b_i will be called a *partial quotient* or *term* in the continued fraction.

We will assume that whenever we discuss a continued fraction $[c, b_0, b_1, \dots, b_n]$, it does, in fact, evaluate to a rational number. For example, $[c, 2, -1, 2]$ is not a valid continued fraction since

$$c + \frac{1}{2 + \frac{1}{-1 + \frac{1}{2}}} = c + \frac{1}{2 + \frac{1}{\frac{-1}{2}}} = c + \frac{1}{2 - 2}$$

is not defined as a rational number.

Conway notation and continued fractions are related in that we can calculate a rational knot's associated continued fraction by reversing the order of the Conway notation. That is, a knot with Conway notation $a_0 a_1 \dots a_n$ has the continued fraction $[0, a_n, a_{n-1}, \dots, a_0] = \frac{p}{q}$. We will denote this knot $K(\frac{p}{q})$. In fact (see Cromwell [2] for a proof), all tangles given by Conway notation corresponding to the same rational number are equivalent. Passing to knots introduces additional equivalences: two rational knots $K(\frac{p}{q})$ and $K(\frac{p'}{q})$ are equivalent if and only if $p' \equiv p^{\pm 1} \pmod{q}$. Also, the knots $K(\frac{p}{q})$ and $K(-\frac{p}{q})$ are mirror reflections of one another.

Recall that if, in a continued fraction $\frac{p}{q} = [c, a_0, \dots, a_n]$, we restrict every a_i , $0 \leq i \leq n$, to be a positive integer, with $a_n > 1$, then this representation of $\frac{p}{q}$ is unique. We will call this the *simple continued fraction* of $\frac{p}{q}$. The corresponding rational knot with Conway notation $a_n \dots a_0$ then has crossing number $c(K) = \sum_{i=0}^n a_i$, as is proven by Ernst and Sumners [3].

Throughout this paper, we will use the notation $(b_0, \dots, b_m)^c$ to mean that the pattern " $b_0, \dots, b_m$ " is repeated c times, with c being any nonnegative integer, e.g., $[0, (-2, 2)^2] = [0, -2, 2, -2, 2]$ and $[0, (-2, 2)^0, 2] = [0, 2]$.

3. BOUNDARY SLOPES

In this section, we will briefly review how to calculate boundary slopes for rational knots.

Let $B(K)$, or simply B , denote the set of all boundary slopes for a knot K . For rational knots $K(\frac{p}{q})$, recall [4] that B can be calculated from continued fractions of $\frac{p}{q}$ having every partial quotient at least two in absolute value. (We will refer to these as *boundary slope continued fractions*.) Specifically, one takes such a continued fraction and pattern-matches the partial quotients against the pattern $[+ - + - \dots]$. The number of terms matching this pattern we call b^+ , and the number of terms not matching this pattern (e.g., the total number of terms minus b^+) we call b^- (since these terms match the pattern $[- + - + \dots]$). In this way, we associate to each boundary slope continued fraction two non-negative integers b^+ and b^- .

Among the boundary slope continued fractions, there is a unique one consisting only of even terms (indeed, this is easy to see using the four substitutions we will derive in Section 5). This corresponds to a Seifert surface of boundary slope 0, so we will denote its b^+ and b^- by b_0^+ and b_0^- . Then, the boundary slope associated to any other continued fraction is given by comparing its b^+ and b^- with those of the Seifert slope; the boundary slope is $2((b^+ - b^-) - (b_0^+ - b_0^-))$. Applying this calculation to every continued fraction with terms at least two in absolute value gives the set of boundary slopes B . B is a finite set of even integers. The *diameter* $D(K)$ is the difference between the maximum and minimum elements of B .

4. CONTINUED FRACTION IDENTITIES

In this section, we will prove four identities related to continued fractions. For identities 2 and 4, we will allow the last entry in a continued fraction to be any nonzero rational number, provided the resulting continued fraction represents a rational number. Note that $[b_0, \dots, b_m, k] = [b_0, \dots, b_m, a_0, \dots, a_n]$ when $k = [a_0, \dots, a_n]$.

Throughout this section, let $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ and $\mathbb{Q}^* = \mathbb{Q} \setminus \{0\}$.

Identity 1. *Let $c \in \mathbb{N}$. Then*

$$[(-2, 2)^c] = -\frac{2c+1}{2c}$$

Proof. By induction.

Base Case ($c = 1$): $[-2, 2] = -2 + \frac{1}{2} = -\frac{3}{2} = -\frac{2 \cdot 1 + 1}{2 \cdot 1}$.

Induction Step: Assume that $[(-2, 2)^c] = \frac{2c+1}{2c}$. Then

$$\begin{aligned} [(-2, 2)^{c+1}] &= -2 + \frac{1}{2 + \frac{1}{[(-2, 2)^c]}} \\ &= -2 + \frac{1}{2 + \frac{1}{-\frac{2c+1}{2c}}} = -2 + \frac{2c+1}{2c+2} \\ &= -\frac{2c+3}{2c+2} = -\frac{2(c+1)+1}{2(c+1)} \end{aligned}$$

■

Identity 2. *Let $c \in \mathbb{N}_0$ and $k \in \mathbb{Q}^*$. Then*

$$[(-2, 2)^c, k] = \frac{2ck + 2c + k}{1 - 2ck - 2c}$$

Note that the denominator becomes zero only in the case where the continued fraction does not represent a rational number.

Proof. By induction.

Base Case ($c = 0$): $[k] = k = \frac{2 \cdot 0 \cdot k + 2 \cdot 0 + k}{1 - 2 \cdot 0 \cdot k - 2 \cdot 0}$.

Induction Step: Assume that $[(-2, 2)^c, k] = \frac{2ck + 2c + k}{1 - 2ck - 2c}$. Then

$$\begin{aligned} [(-2, 2)^{c+1}, k] &= -2 + \frac{1}{2 + \frac{1}{[(-2, 2)^c, k]}} \\ &= -2 + \frac{1}{2 + \frac{1}{\frac{2ck + 2c + k}{1 - 2ck - 2c}}} \\ &= -2 + \frac{2ck + 2c + k}{2ck + 2c + 2k + 1} \\ &= \frac{-2ck - 2c - 3k - 2}{2ck + 2c + 2k + 1} \\ &= \frac{2(c+1)k + 2(c+1) + k}{1 - 2(c+1)k - 2(c+1)} \end{aligned}$$

■

Identity 3. *Let $c \in \mathbb{N}$. Then*

$$[(2, -2)^c] = \frac{2c+1}{2c}$$

Proof. Note that $[(2, -2)^c] = [2, (-2, 2)^{c-1}, -2]$. Apply Identity 2. ■

Identity 4. *Let $c \in \mathbb{N}_0$ and $k \in \mathbb{Q}^*$. Then*

$$[(2, -2)^c, k] = \frac{2ck - 2c + k}{2ck - 2c + 1}$$

Again, the denominator becomes zero only in the case where the continued fraction does not represent a rational number.

Proof. This proof will be done in two parts.

Case 1 ($c = 0$): $[k] = k = \frac{2 \cdot 0 \cdot k - 2 \cdot 0 + k}{2 \cdot 0 \cdot k - 2 \cdot 0 + 1}$.

Case 2 ($c > 0$): Note that $[(2, -2)^c, k] = [2, (-2, 2)^{c-1}, -2, k]$. Apply Identity 2. ■

5. SUBSTITUTION RULES

In this section, we will prove four identities, or substitutions, which will be used to derive equal continued fractions. In particular, given $\frac{p}{q} = [c, a_0, \dots, a_n]$, where every a_i is positive, these substitutions can be used to calculate all the boundary slope continued fractions of $\frac{p}{q}$, i.e., continued fractions $[c', b_0, \dots, b_m]$ where each $|b_i| \geq 2$. We conclude the section with an example to illustrate how these rules can be applied to a specific continued fraction.

Throughout this section, let $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ and $\mathbb{Z}^* = \mathbb{Z} \setminus \{0\}$.

Substitution 1. *Let $n \in \mathbb{N}$. Let $a_0 \in \mathbb{Z}$ and $a_1 \in \mathbb{N}$. If $n = 2$ then let $a_2 \in \mathbb{Z} \setminus \{0, -1\}$. If $n \geq 3$ then let $a_i \in \mathbb{Z}^*$ for all $2 \leq i \leq n$. If we have $r = [a_0, 2a_1, a_2, a_3, \dots, a_n]$, then $r = [a_0 + 1, (-2, 2)^{a_1-1}, -2, a_2 + 1, a_3, a_4, \dots, a_n]$.*

Proof. This proof will be done in three parts.

Case 1 ($n = 1$): We want to show that $[a_0, 2a_1] = [a_0 + 1, (-2, 2)^{a_1-1}, -2]$.

$$\begin{aligned} & [a_0 + 1, (-2, 2)^{a_1-1}, -2] \\ = & a_0 + 1 + \frac{1}{[(-2, 2)^{a_1-1}, -2]} \\ = & a_0 + 1 + \frac{1}{\frac{2(a_1-1)(-2) + 2(a_1-1) + (-2)}{1 - 2(a_1-1)(-2) - 2(a_1-1)}} \quad (\text{Apply Identity 2}) \\ = & a_0 + 1 + \frac{-2a_1 + 1}{2a_1} \\ = & a_0 + \frac{1}{2a_1} \\ = & [a_0, 2a_1] \end{aligned}$$

Case 2 ($n = 2$): We want to show $[a_0, 2a_1, a_2] = [a_0 + 1, (-2, 2)^{a_1-1}, -2, a_2 + 1]$.

$$\begin{aligned}
& [a_0 + 1, (-2, 2)^{a_1-1}, -2, a_2 + 1] \\
= & a_0 + 1 + \frac{1}{[(-2, 2)^{a_1-1}, -2, a_2 + 1]} \\
= & a_0 + 1 + \frac{1}{\frac{2(a_1-1)\left(\frac{-2a_2-1}{a_2+1}\right) + 2(a_1-1) + \frac{-2a_2-1}{a_2+1}}{1-2(a_1-1)\left(\frac{-2a_2-1}{a_2+1}\right) - 2(a_1-1)}} \quad (\text{Apply Identity 2}) \\
= & a_0 + 1 + \frac{a_2 - 2a_1a_2 - 1}{2a_1a_2 + 1} \\
= & a_0 + \frac{1}{\frac{2a_1a_2+1}{a_2}} \\
= & a_0 + \frac{1}{2a_1 + \frac{1}{a_2}} \\
= & [a_0, 2a_1, a_2]
\end{aligned}$$

Case 3 ($n \geq 3$): We want to show that $[a_0, 2a_1, a_2, a_3, a_4, \dots, a_n] = [a_0 + 1, (-2, 2)^{a_1-1}, -2, a_2 + 1, a_3, a_4, \dots, a_n]$. Let $R = [a_3, a_4, \dots, a_n]$.

$$\begin{aligned}
& [a_0 + 1, (-2, 2)^{a_1-1}, -2, a_2 + 1, a_3, a_4, \dots, a_n] \\
= & a_0 + 1 + \frac{1}{[(-2, 2)^{a_1-1}, -2, a_2 + 1, a_3, a_4, \dots, a_n]} \\
= & a_0 + 1 + \frac{1}{\frac{2(a_1-1)\left(\frac{-2Ra_2-R-2}{Ra_2+R+1}\right) + 2(a_1-1) + \frac{-2Ra_2-R-2}{Ra_2+R+1}}{1-2(a_1-1)\left(\frac{-2Ra_2-R-2}{Ra_2+R+1}\right) - 2(a_1-1)}} \quad (\text{Apply Identity 2}) \\
= & a_0 + 1 + \frac{Ra_2 - 2a_1 - R - 2Ra_1a_2 + 1}{R + 2a_1 + 2Ra_1a_2} \\
= & a_0 + \frac{1}{\frac{R+2a_1+2Ra_1a_2}{Ra_2+1}} \\
= & a_0 + \frac{1}{2a_1 + \frac{R}{Ra_2+1}} \\
= & a_0 + \frac{1}{2a_1 + \frac{1}{\frac{1}{a_2} + \frac{1}{R}}} \\
= & [a_0, 2a_1, a_2, a_3, a_4, \dots, a_n]
\end{aligned}$$

■

Substitution 2. Let $n \in \mathbb{N}$. Let $a_0 \in \mathbb{Z}$ and $a_1 \in \mathbb{N}$. If $n = 2$ then let $a_2 \in \mathbb{Z} \setminus \{0, 1\}$. If $n \geq 3$ then let $a_i \in \mathbb{Z}^*$ for all $2 \leq i \leq n$. If we have $r = [a_0, -2a_1, a_2, a_3, \dots, a_n]$, then $r = [a_0 - 1, (2, -2)^{a_1-1}, 2, a_2 - 1, a_3, a_4, \dots, a_n]$.

Proof. This proof will be done in three parts.

Case 1 ($n = 1$): We want to show that $[a_0, -2a_1] = [a_0 - 1, (2, -2)^{a_1-1}, 2]$.

$$\begin{aligned}
[a_0 - 1, (2, -2)^{a_1-1}, 2] &= a_0 - 1 + \frac{1}{[(2, -2)^{a_1-1}, 2]} \\
&= a_0 - 1 + \frac{1}{\frac{2(a_1-1)(2) - 2(a_1-1) + (2)}{2(a_1-1)(2) - 2(a_1-1) + 1}} \quad (\text{Apply Identity 4}) \\
&= a_0 - 1 + \frac{2a_1 - 1}{2a_1} \\
&= a_0 + \frac{1}{-2a_1} \\
&= [a_0, -2a_1]
\end{aligned}$$

Case 2 ($n = 2$): We want to show $[a_0, -2a_1, a_2] = [a_0 - 1, (2, -2)^{a_1-1}, 2, a_2 - 1]$.

$$\begin{aligned}
&[a_0 - 1, (2, -2)^{a_1-1}, 2, a_2 - 1] \\
&= a_0 - 1 + \frac{1}{[(2, -2)^{a_1-1}, 2, a_2 - 1]} \\
&= a_0 - 1 + \frac{1}{\frac{2(a_1-1)\left(\frac{2a_2-1}{a_2-1}\right) - 2(a_1-1) + \frac{2a_2-1}{a_2-1}}{2(a_1-1)\left(\frac{2a_2-1}{a_2-1}\right) - 2(a_1-1) + 1}} \quad (\text{Apply Identity 4}) \\
&= a_0 - 1 + \frac{2a_1a_2 - a_2 - 1}{2a_1a_2 - 1} \\
&= a_0 + \frac{1}{\frac{2a_1a_2-1}{-a_2}} \\
&= a_0 + \frac{1}{-2a_1 + \frac{1}{a_2}} \\
&= [a_0, -2a_1, a_2]
\end{aligned}$$

Case 3 ($n \geq 3$): We want to show that $[a_0, -2a_1, a_2, a_3, a_4, \dots, a_n] = [a_0 - 1, (2, -2)^{a_1-1}, 2, a_2 - 1, a_3, a_4, \dots, a_n]$. Let $R = [a_3, a_4, \dots, a_n]$.

$$\begin{aligned}
&[a_0 - 1, (2, -2)^{a_1-1}, 2, a_2 - 1, a_3, a_4, \dots, a_n] \\
&= a_0 - 1 + \frac{1}{[(2, -2)^{a_1-1}, 2, a_2 - 1, a_3, a_4, \dots, a_n]} \\
&= a_0 - 1 + \frac{1}{\frac{2(a_1-1)\left(\frac{2Ra_2+2-R}{Ra_2+1-R}\right) - 2(a_1-1) + \frac{2Ra_2+2-R}{Ra_2+1-R}}{2(a_1-1)\left(\frac{2Ra_2+2-R}{Ra_2+1-R}\right) - 2(a_1-1) + 1}} \quad (\text{Apply Identity 4}) \\
&= a_0 - 1 + \frac{2a_1 - R - Ra_2 + 2Ra_1a_2 - 1}{2a_1 - R + 2Ra_1a_2} \\
&= a_0 + \frac{1}{\frac{R-2a_1-2Ra_1a_2}{Ra_2+1}} \\
&= a_0 + \frac{1}{-2a_1 + \frac{R}{Ra_2+1}} \\
&= a_0 + \frac{1}{-2a_1 + \frac{1}{a_2 + \frac{1}{R}}} \\
&= [a_0, -2a_1, a_2, a_3, a_4, \dots, a_n]
\end{aligned}$$

■

Substitution 3. Let $n \in \mathbb{N}$. Let $a_0 \in \mathbb{Z}$ and $a_1 \in \mathbb{N}_0$. If $n = 2$ then let $a_2 \in \mathbb{Z} \setminus \{0, -1\}$. If $n \geq 3$ then let $a_i \in \mathbb{Z}^*$ for all $2 \leq i \leq n$. If we have $r = [a_0, 2a_1 + 1, a_2, a_3, \dots, a_n]$, then $r = [a_0 + 1, (-2, 2)^{a_1}, -a_2 - 1, -a_3, -a_4, \dots, -a_n]$.

Proof. This proof will be done in three parts.

Case 1 ($n = 1$): We want to show that $[a_0, 2a_1 + 1] = [a_0 + 1, (-2, 2)^{a_1}]$. Note: When $a_1 = 0$ this is trivially true. So, we can assume $a_1 > 0$.

$$\begin{aligned}
 [a_0 + 1, (-2, 2)^{a_1}] &= a_0 + 1 + \frac{1}{[(-2, 2)^{a_1}]} \\
 &= a_0 + 1 + \frac{1}{-\frac{2a_1+1}{2a_1}} \quad (\text{Apply Identity 1}) \\
 &= a_0 + 1 + \frac{2a_1}{-2a_1 - 1} \\
 &= a_0 + \frac{1}{2a_1 + 1} \\
 &= [a_0, 2a_1 + 1]
 \end{aligned}$$

Case 2 ($n = 2$): We want to show $[a_0, 2a_1 + 1, a_2] = [a_0 + 1, (-2, 2)^{a_1}, -a_2 - 1]$.

$$\begin{aligned}
 &[a_0 + 1, (-2, 2)^{a_1}, -a_2 - 1] \\
 &= a_0 + 1 + \frac{1}{[(-2, 2)^{a_1}, -a_2 - 1]} \\
 &= a_0 + 1 + \frac{1}{\frac{2a_1(-a_2-1)+2a_1+(-a_2-1)}{1-2a_1(-a_2-1)-2a_1}} \quad (\text{Apply Identity 2}) \\
 &= a_0 + 1 + \frac{-2a_1a_2 - 1}{2a_1a_2 + a_2 + 1} \\
 &= a_0 + \frac{1}{\frac{2a_1a_2+a_2+1}{a_2}} \\
 &= a_0 + \frac{1}{2a_1 + 1 + \frac{1}{a_2}} \\
 &= [a_0, 2a_1 + 1, a_2]
 \end{aligned}$$

Case 3 ($n \geq 3$): We want to show that $[a_0, 2a_1 + 1, a_2, a_3, a_4, \dots, a_n] = [a_0 + 1, (-2, 2)^{a_1}, -a_2 - 1, -a_3, -a_4, \dots, -a_n]$. Let $R = [a_3, a_4, \dots, a_n]$. Note: $-R =$

$$[-a_3, -a_4, \dots, -a_n].$$

$$\begin{aligned}
& [a_0 + 1, (-2, 2)^{a_1}, -a_2 - 1, -a_3, -a_4, \dots, -a_n] \\
= & a_0 + 1 + \frac{1}{[(-2, 2)^{a_1}, -a_2 - 1, -a_3, -a_4, \dots, -a_n]} \\
= & a_0 + 1 + \frac{1}{\frac{2a_1 \left(\frac{Ra_2 + R + 1}{-R} \right) + 2a_1 + \frac{Ra_2 + R + 1}{-R}}{1 - 2a_1 \left(\frac{Ra_2 + R + 1}{-R} \right) - 2a_1}} \quad (\text{Apply Identity 2}) \\
= & a_0 + 1 + \frac{-R - 2a_1 - 2Ra_1a_2}{2a_1 + R + Ra_2 + 2Ra_1a_2 + 1} \\
= & a_0 + \frac{1}{\frac{2a_1 + R + Ra_2 + 2Ra_1a_2 + 1}{1 + Ra_2}} \\
= & a_0 + \frac{1}{2a_1 + 1 + \frac{1}{\frac{Ra_2 + 1}{R}}} \\
= & a_0 + \frac{1}{2a_1 + 1 + \frac{1}{a_2 + \frac{1}{R}}} \\
= & [a_0, 2a_1 + 1, a_2, a_3, a_4, \dots, a_n]
\end{aligned}$$

■

Substitution 4. Let $n \in \mathbb{N}$. Let $a_0 \in \mathbb{Z}$ and $a_1 \in \mathbb{N}_0$. If $n = 2$ then let $a_2 \in \mathbb{Z} \setminus \{0, 1\}$. If $n \geq 3$ then let $a_i \in \mathbb{Z}^*$ for all $2 \leq i \leq n$. If we have $r = [a_0, -2a_1 - 1, a_2, a_3, \dots, a_n]$, then $r = [a_0 - 1, (2, -2)^{a_1}, -a_2 + 1, -a_3, -a_4, \dots, -a_n]$.

Proof. This proof will be done in three parts.

Case 1 ($n = 1$): We want to show that $[a_0, -2a_1 - 1] = [a_0 - 1, (2, -2)^{a_1}]$. Note: When $a_1 = 0$ this is trivially true. So, we can assume $a_1 > 0$.

$$\begin{aligned}
[a_0 - 1, (2, -2)^{a_1}] &= a_0 - 1 + \frac{1}{[(2, -2)^{a_1}]} \\
&= a_0 - 1 + \frac{1}{\frac{2a_1 + 1}{2a_1}} \quad (\text{Apply Identity 3}) \\
&= a_0 - 1 + \frac{2a_1}{2a_1 + 1} \\
&= a_0 + \frac{1}{-2a_1 - 1} \\
&= [a_0, -2a_1 - 1]
\end{aligned}$$

Case 2 ($n = 2$): We want to show $[a_0, -2a_1 - 1, a_2] = [a_0 - 1, (2, -2)^{a_1}, -a_2 + 1]$.

$$\begin{aligned}
& [a_0 - 1, (2, -2)^{a_1}, -a_2 + 1] \\
&= a_0 - 1 + \frac{1}{[(2, -2)^{a_1}, -a_2 + 1]} \\
&= a_0 - 1 + \frac{1}{\frac{2a_1(-a_2+1) - 2a_1 + (-a_2+1)}{2a_1(-a_2+1) - 2a_1 + 1}} \quad (\text{Apply Identity 4}) \\
&= a_0 - 1 + \frac{2a_1a_2 - 1}{2a_1a_2 + a_2 - 1} \\
&= a_0 + \frac{1}{\frac{1-a_2-2a_1a_2}{a_2}} \\
&= a_0 + \frac{1}{-2a_1 - 1 + \frac{1}{a_2}} \\
&= [a_0, -2a_1 - 1, a_2]
\end{aligned}$$

Case 3 ($n \geq 3$): We want to show that $[a_0, -2a_1 - 1, a_2, a_3, a_4, \dots, a_n] = [a_0 - 1, (2, -2)^{a_1}, -a_2 + 1, -a_3, -a_4, \dots, -a_n]$. Let $R = [a_3, a_4, \dots, a_n]$. Note: $-R = [-a_3, -a_4, \dots, -a_n]$.

$$\begin{aligned}
& [a_0 - 1, (2, -2)^{a_1}, -a_2 + 1, -a_3, -a_4, \dots, -a_n] \\
&= a_0 - 1 + \frac{1}{[(2, -2)^{a_1}, -a_2 + 1, -a_3, -a_4, \dots, -a_n]} \\
&= a_0 - 1 + \frac{1}{\frac{2a_1\left(\frac{R-Ra_2-1}{R}\right) - 2a_1 + \frac{R-Ra_2-1}{R}}{2a_1\left(\frac{R-Ra_2-1}{R}\right) - 2a_1 + 1}} \quad (\text{Apply Identity 4}) \\
&= a_0 - 1 + \frac{2a_1 - R + 2Ra_1a_2}{2a_1 - R + Ra_2 + 2Ra_1a_2 + 1} \\
&= a_0 + \frac{1}{\frac{R-2a_1-Ra_2-2Ra_1a_2-1}{Ra_2+1}} \\
&= a_0 + \frac{1}{-2a_1 - 1 + \frac{1}{\frac{Ra_2+1}{R}}} \\
&= a_0 + \frac{1}{-2a_1 - 1 + \frac{1}{a_2 + \frac{1}{R}}} \\
&= [a_0, -2a_1 - 1, a_2, a_3, a_4, \dots, a_n]
\end{aligned}$$

■

5.1. An example of the application of Substitutions 1–4. Let us illustrate how the above results can be used to generate a list of all boundary slope continued fractions starting from the simple continued fraction. As an example, suppose we start with $[0, 2a, 2b+1, 2c]$, where $a, c \in \mathbb{N}$ and $b \in \mathbb{N} \cup \{0\}$. By applying Substitution 1, we can immediately derive another continued fraction: $[1, (-2, 2)^{a-1}, -2, 2b+2, 2c]$. We will refer to this as *applying Substitution 1 at position 0* as it is the a_0 term, $2a$, that has been replaced by the sequence $-2, 2, \dots, -2$.

Applying the same substitution at position 2, we get $[1, (-2, 2)^{a-1}, -2, 2b+3, (-2, 2)^{c-1}, -2]$. We could continue on this path, but it is easy to see that any further substitutions will result in a ± 1 term. Therefore, we return to the original

sequence and use Substitution 3 (at position 1) to obtain $[0, 2a+1, (-2, 2)^b, -2c-1]$. Finally, applying Substitution 1 at position 2, we have $[0, 2a, 2b+2, (-2, 2)^{c-1}, -2]$.

Thus, there are five boundary slope continued fractions that can be derived from the simple continued fraction $[0, 2a, 2b+1, 2c]$: three obtained by substitutions at positions 0, 1, and 2; one by substitutions at 0 and 2; and the original continued fraction itself (with no substitutions). Note, that these are precisely the fractions obtained by applying substitutions at non-adjacent positions.

6. PROOF OF THEOREM 2

In this section we will prove Theorem 2, that the boundary slope continued fractions are among the fractions obtained by applying substitutions at non-adjacent positions in the original simple continued fraction. Our strategy is to first review Langford's argument [8] that the boundary slopes are determined by the leaves of a binary tree. We then show, by induction, that applying substitutions at non-adjacent positions accounts for all the leaves of the tree.

6.1. The boundary slope binary tree. Recall that for any rational $\frac{p}{q}$, we can find another rational $\frac{p'}{q}$ such that $0 \leq \frac{p'}{q} < 1$ and $K\left(\frac{p}{q}\right) = K\left(\frac{p'}{q}\right)$. Also, recall that the rational 0 corresponds to the unknot, which has a rather boring set of continued fractions (namely, 0 is the only one). Therefore, without loss of generality, we will assume henceforth that $0 < \frac{p}{q} < 1$. There is a unique simple continued fraction, $[0, a_0, \dots, a_n]$, for such $\frac{p}{q}$, such that $a_n \geq 2$ and, for all $i \in \{0, 1, \dots, n\}$, $a_i > 0$.

Before we can prove Theorem 2, we must first state a lemma. The straightforward proof by induction may be found in Langford [8] which is also the source for the following definition.

Definition 1. *The k th subexpansion of $[c, a_0, \dots, a_n]$ is the continued fraction $[0, a_k, \dots, a_n]$ where $0 \leq k \leq n$.*

Lemma 1. *Let $[c, a_0, \dots, a_n]$ be a boundary slope continued fraction, that is, for each $i \in \{0, 1, \dots, n\}$, $|a_i| \geq 2$. Then every subexpansion r of $[c, a_0, \dots, a_n]$ satisfies $|r| < 1$.*

As Langford [8] has shown, a complete list of boundary slope continued fractions for a rational $\frac{p}{q}$, where each partial quotient is at least two in absolute value, can be calculated by means of a binary tree. We will now outline the creation of this binary tree which follows from Lemma 1.

The root vertex is labelled with the fraction $\frac{p}{q}$ and the two edges coming from the root are labelled $0 = \lfloor \frac{p}{q} \rfloor$ and $1 = \lceil \frac{p}{q} \rceil$. At every other vertex in the tree, we arrive with the first k terms in a continued fraction for $\frac{p}{q}$ and a rational number r representing the $(k-1)$ st subexpansion. The k terms are found as labels of the edges of the tree starting from the root and continuing to the vertex in question. We label the vertex with r . Since, by Lemma 1, any k th subexpansion is less than one in absolute value, we know that the next term in the continued fraction, a_i , is within 1 of $1/r$: $|a_i - 1/r| < 1$. However, a_i is an integer. Therefore, a_i is either the floor $\lfloor 1/r \rfloor$ or the ceiling $\lceil 1/r \rceil$ of $1/r$. If $1/r$ is not an integer, there will be two edges coming out of the vertex, one labelled with $\lfloor 1/r \rfloor$, and the other labelled with

$\lceil 1/r \rceil$. Since $|r| < 1$, neither of these arrows is 0. If either is ± 1 , we terminate that edge with a leaf labelled “ $\nexists$ ” to indicate that this path does not lead to a boundary slope continued fraction. (When we refer to the leaves of the binary tree below, we will be excluding these “dead” leaves.) If $1/r$ is an integer, then, there is only one edge coming out of the vertex. Label the edge with $1/r$ and label the leaf vertex at the end of this edge with the continued fraction expansion for $\frac{p}{q}$ given by the labels of the edges from the root to the leaf.

For example, Figure 1 shows the binary tree for the fraction $2/7$ (which corre-




FIGURE 1. The boundary slope binary tree for $\frac{p}{q} = \frac{2}{7}$ (the 5_2 knot).

sponds to the 5_2 knot).

Thus, by Lemma 1, the algorithm used to construct the tree will provide all the boundary slope continued fractions of $\frac{p}{q}$ as leaf vertices.

6.2. Binary tree from substitutions. Now, let’s prove the theorem by showing that the leaves of Langford’s binary tree (and therefore the set of boundary slopes) correspond to applying substitutions at non-adjacent positions in the simple continued fraction.

Theorem 2. *The boundary slope continued fractions of $K(p/q)$ are among the continued fractions obtained by applying substitutions at non-adjacent positions in the simple continued fraction of p/q .*

Proof. We proceed by induction on the length n of the simple continued fraction $[0, a_0, a_1, \dots, a_n]$.

Case 1 ($n = 0$): Here, $p/q = 1/a_0$. We wish to show that the boundary slope continued fractions are among the two continued fractions given by substituting or not at position 0. There are three subcases. (To simplify the exposition, we will not be considering the, very similar, trees that arise when the terms a_i are negative although they may be required as part of our induction.)

Subcase 1 ($a_0 = 1$): In this case, the tree is shown in Figure 2. There are no




FIGURE 2. The binary tree for $[0, 1]$.

boundary slope continued fractions in this case. (Actually, here $\frac{p}{q} = 1$, so we've violated our assumption that $\frac{p}{q} < 1$. Ordinarily, we would represent this knot, the unknot, by $[0]$ and that would also be the only boundary slope. We include this case as it may arise as part of our induction.) Thus, it is true that all boundary slope continued fractions are among the two continued fractions $[0, 1]$ and $[1]$ given by substituting or not at position 0.

Subcase 2 ($a_0 = 2a$, $a \geq 1$): The binary tree is shown in Figure 3. There are two boundary slope continued fractions, and they are the fractions $[0, a_0]$ and $[1, (-2, 2)^a, -2]$ given by substituting or not at position 0.

Subcase 3 ($a_0 = 2a + 1$, $a \geq 1$): The binary tree is shown in Figure 4. The two boundary slope continued fractions $[0, a_0]$ and $[1, (-2, 2)^a]$ are those given by substituting or not at position 0.


Case 2 ($n = 1$): Our goal is to show that the boundary slope continued fractions are among the fractions given by substituting at position 0, at position 1, and by not substituting at all. The result of substitution at position 0 will depend on whether a_0 is even or odd:

$$\begin{aligned} [0, 2a, a_1] &\xrightarrow{\text{Sub. } 1} [1, (-2, 2)^{(a-1)}, -2, a_1 + 1] \\ [0, 2a + 1, a_1] &\xrightarrow{\text{Sub. } 3} [1, (-2, 2)^a, -a_1 - 1] \end{aligned}$$

Similarly, substitution at position 1 depends on the parity of a_1 :


$$\begin{aligned} [0, a_0, 2b] &\xrightarrow{\text{Sub. } 1} [0, a_0 + 1, (-2, 2)^{(b-1)}, -2] \\ [0, a_0, 2b + 1] &\xrightarrow{\text{Sub. } 3} [0, a_0 + 1, (-2, 2)^b] \end{aligned}$$

As Figure 5 shows, these two boundary slopes, along with the original continued fraction $[0, a_0, a_1]$ (no substitutions) are precisely those that arise in the binary tree. Note that if, for example, a_0 or a_1 is 1, then the $[0, a_0, a_1]$ leaf is not in fact a boundary slope continued fraction. The point is that all leaves of the binary tree are included in the set of continued fractions obtained by substitutions at non-adjacent positions. So, every boundary slope continued fraction appears in this set.

FIGURE 3. The binary tree for $[0, 2a]$.


Case 3 ($n = 2$): This case will illustrate how the induction works. There are five continued fractions given by substitutions at non-adjacent positions (compare with the example of Section 5.1): three obtained by substitutions at positions 0, 1, and 2; one by substitutions at 0 and 2; and the original continued fraction itself (with no substitutions). Let us denote these choices of substitutions by a sequence of three 0's and 1's where a 1 in the i th place denotes a substitution at that i th position. Thus, the five continued fractions will be denoted 100, 010, 001, 101, and 000.

We can think of the binary tree (Figure 6) as being a union of two subtrees. The one at left corresponds to making no substitution at position 0. This subtree ends in the three boundary slopes which have: no substitutions (000); substitution at

FIGURE 4. The binary tree for $[0, 2a + 1]$.


position 1 (010); and substitution at position 2 (001), i.e., the sequences that begin in 0. This subtree is essentially the same as that for the $[0, a_1, a_2]$ continued fraction (compare Figure 5) as we can obtain these three sequences by adding a 0 at the front of the three boundary slopes sequences 00, 10, and 01 of that case. The other subtree corresponds to making a substitution at position 0 and no substitution at position 1. This subtree contains the remaining two boundary slopes: substitution at position 0 (100); and substitution at positions 0 and 2 (101), i.e., sequences that begin in 10. This subtree is similar to that for $[0, a_2]$ (compare Figure 3) as it remains only to decide whether or not to substitute in the second position. Again, some of these five sequences may not result in a boundary slope continued fraction, for example, if one of the a_i is 1. However, every leaf of the tree will be included in the set of continued fractions obtained by substituting at non-adjacent positions.

Case 4 ($n \geq 3$): As in Case 3, we can decompose the binary tree (Figure 7) into two subtrees. One corresponds to sequences that begin with 0, the other to

FIGURE 5. The binary tree for $[0, a_0, a_1]$.

sequences beginning with 10. The first will be, essentially, the tree that arises from the simple continued fraction $[0, a_1, a_2, \dots, a_n]$. By induction, the leaves of this subtree correspond to non-adjacent substitutions in this simple continued fraction. By its placement in the $[0, a_0, a_1, \dots, a_n]$ tree, this ensures that the leaves of this part of the tree will correspond to continued fractions obtained by substitution sequences into $[0, a_0, a_1, \dots, a_n]$ that begin with 0.

The other subtree is isomorphic to the tree that arises from the simple continued fraction $[0, a_2, a_3, \dots, a_n]$. By induction, the leaves of the subtree correspond to substitutions into this continued fraction. By its placement in the tree


 FIGURE 6. The $[0, a_0, a_1, a_2]$ tree is a union of two subtrees.

for $[0, a_0, a_1, \dots, a_n]$, the leaves here can be obtained by non-adjacent substitutions into that continued fraction that begin with 10.

Thus, every leaf of the binary tree and, therefore, every boundary slope continued fraction can be obtained by non-adjacent substitutions into the simple continued fraction. $\blacksquare$




FIGURE 7. The general case also results in two subtrees.

Corollary 2. *If $\frac{p}{q} = [0, a_0, a_1, \dots, a_n]$ is a simple continued fraction, then $K(p/q)$ has at most F_{n+2} boundary slopes where F_n is the n th Fibonacci number.*

Proof. We have shown that the boundary slope continued fractions lie among those given by substitution at non-adjacent positions which in turn are in bijection with sequences of $n+1$ 0's or 1's containing no pair of consecutive 1's. Thus the number of boundary slopes is at most P_n , where P_n is the number of 0, 1 sequences of length $n+1$ with no consecutive 1's. We will show that $P_n = F_{n+2}$ by induction.

There are two base cases. If $n = 0$, there are two sequences: 0 and 1. So, $P_0 = 2 = F_2$. For $n = 1$, there are three sequences: 00, 10, and 01. So, $P_1 = 3 = F_3$.

For the inductive step, sequences of length $n+1$ are obtained by either adding a 0 to the beginning of a n sequence or 10 to the beginning of a $n-1$ sequence. Thus $P_n = P_{n-1} + P_{n-2} = F_{n+1} + F_n = F_{n+2}$. ■

In general, F_{n+2} is an overestimate since the continued fractions obtained by substitutions will not necessarily have terms at least two in absolute value. In particular, if the simple continued fraction includes any 1's, then the continued fraction obtained by making no substitutions (000...0) will not be a boundary slope continued fraction. Moreover, different boundary slope continued fractions could result in the same boundary slope. For example, this will occur when, in the simple continued fraction, we have two equal terms separated by an even distance: $a_i = a_{i+2k}$.

7. MAXIMUM AND MINIMUM BOUNDARY SLOPES

In this section, we will show how one can calculate the maximum and minimum boundary slopes. We will refer to the minimum value of $b^+ - b^-$ as b_1 and the maximum as b_2 . Further, we will refer to the components of b_1 as b_1^+ and b_1^- . Similarly for b_2 .

The key observation is that, if we begin with a simple continued fraction, applying a substitution at an even position will decrease $b^+ - b^-$ and, hence, the boundary slope, while applying a substitution at an odd position will increase the boundary slope. Thus, we can minimize the boundary slope by substituting at each even position (and no odd positions). Note that this will result in a continued fraction where each term is at least two in absolute value. Indeed, the even position terms of the original simple continued fraction will be replaced by a sequence of ± 2 's while the terms in the odd positions will be augmented in absolute value by at least one.

However, we need a way to count the resulting b_1^+ and b_1^- when we make substitutions at each even position. We will do this by focusing on the clusters of $(\pm 2, \mp 2)$, examining what occurs near them individually, and then summing up the results. We replace an even number, $2k$, with $(\pm 2, \mp 2)^{|k|-1}, \pm 2$, or $2|k| - 1$ terms. For a positive odd number $2k + 1$, we replace it with $(\pm 2, \mp 2)^k$ or $2k$ terms and a negative odd $-(2k + 1)$ is also replaced with $2k$ terms. We can combine these cases by observing that a term a_i is replaced with $|a_i| - 1$ terms. If we think of making the substitutions at position $0, 2, 4, \dots$ in turn, then, each substitution will not affect the magnitude of later a_{2i} but may change their signs. Thus, if we begin with the simple continued fraction $[0, a_0, \dots, a_n]$, each a_{2i} will be replaced by $a_{2i} - 1$ terms.

All we need now, then, is to count the number of terms that appear between the strings of ± 2 's. There will be one such term between every string of ± 2 's, and possibly one at the tail end of the continued fraction. Specifically, for the simple continued fraction $[0, a_0, \dots, a_n]$ where n is even, there are $\frac{n}{2}$ odd position terms, $a_1, a_3, \dots, a_{n-1}$. Similarly, if n is odd, then we must have $\frac{n+1}{2}$ odd terms. More concisely, there are $\lceil \frac{n}{2} \rceil$ odd terms. Now, we have all the ingredients necessary to calculate b_1^- .

$$\begin{aligned}
 b_1^- &= \left\lceil \frac{n}{2} \right\rceil + \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} (a_{2i} - 1) \\
 &= \left\lceil \frac{n}{2} \right\rceil - \left(\left\lfloor \frac{n}{2} \right\rfloor + 1 \right) + \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} a_{2i} \\
 &= \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} a_{2i} \quad \text{if } n \text{ is odd} \\
 &= -1 + \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} a_{2i} \quad \text{if } n \text{ is even}
 \end{aligned}$$

So, surprisingly, b_1^- is equal to the sum of the even terms, less one if n is even. Recall that, the simple continued fraction is $[0, a_0, \dots, a_n]$ so that n is one more than the number of partial quotients. Note that $b_1^+ = 0$ as all partial quotients are now matched to the $[- + - + \dots]$ pattern.

Similarly, the maximum boundary slope is given by substituting at each odd position, and we can use the exact same logic to find b_2^+ . In fact, the formula itself is nearly identical. The sole differences are the terms we sum over, and the fact that we don't add $\lfloor \frac{n}{2} \rfloor$, but instead $\lceil \frac{n+1}{2} \rceil$. We then get:

$$\begin{aligned}
 b_2^+ &= \left\lceil \frac{n+1}{2} \right\rceil + \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} (a_{2i+1} - 1) \\
 &= \left\lceil \frac{n+1}{2} \right\rceil - \left(\left\lfloor \frac{n-1}{2} \right\rfloor + 1 \right) + \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} a_{2i+1} \\
 &= \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} a_{2i+1} \quad \text{if } n \text{ is odd} \\
 &= 1 + \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} a_{2i+1} \quad \text{if } n \text{ is even}
 \end{aligned}$$

Now that we have b_1 (which is $-b_1^-$, since $b_1^+ = 0$) and b_2 (which is simply b_2^+), we can calculate the maximum and minimum boundary slopes in terms of b_0^+ and b_0^- . The minimum boundary slope is $2((0 - b_1^-) - (b_0^+ - b_0^-)) = -2b_1^- - 2(b_0^+ - b_0^-)$. Similarly, the maximum is $2((b_2^+ - 0) - (b_0^+ - b_0^-)) = 2b_2^+ - 2(b_0^+ - b_0^-)$.

8. PROOF OF THEOREM 1

In this section we prove our main theorem, that twice the crossing number of a 2-bridge knot K is equal to the diameter of the boundary slopes.

Theorem 1. *For K a 2-bridge knot, $D(K) = 2c(K)$.*

Proof. Firstly, calculating the crossing number of a rational knot is simple. If $[0, a_0, \dots, a_n] = \frac{p}{q}$ is the simple continued fraction for $K\left(\frac{p}{q}\right)$, then $c\left(K\left(\frac{p}{q}\right)\right) = \sum_{i=0}^n a_i$.

The diameter of $B(K)$ is also easy to calculate. If we use the b_1 and b_2 from the previous section, we get $D(K) = 2b_2^+ - 2(b_0^+ - b_0^-) - (-2b_1^- - 2(b_0^+ - b_0^-)) = 2b_2^+ + 2b_1^-$. At this point, b_1^- and b_2^+ may vary depending on whether n is even or odd. However, the differences cancel each other out in either instance, leaving us with

$$\begin{aligned}
 D(K) &= 2 \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} a_{2i+1} + 2 \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} a_{2i} \\
 &= 2 \sum_{i=0}^n a_i
 \end{aligned}$$

This concludes the proof that $2c(K) = D(K)$. ■

REFERENCES

- [1] C.C. Adams. *The Knot Book: An Elementary Introduction to the Mathematical Theory of Knots*. American Mathematical Society (2004).
- [2] P. Cromwell. *Knots and Links*. Cambridge University Press (2004).
- [3] C. Ernst and D.W. Sumners. The growth of the number of prime knots. *Math. Proc. Cambridge Phil. Soc.* **102** (1987) 303–315.
- [4] A. Hatcher and W. Thurston. Incompressible surfaces in 2-bridge knot complements. *Inventiones Math.* **79** (1985) 225–246.
- [5] K. Ichihara, Private Communication.
- [6] K. Ichihara and S. Mizushima. Crossing number and diameter of boundary slope set of Montesinos knot. Preprint (2005). math.GT/0510370
- [7] M. Ishikawa and K. Shimokawa. Boundary slopes and crossing numbers of knots. Preprint (2002).
- [8] R. Langford. M.S. Thesis, CSU, Chico (2005).
- [9] L. Moser. Elementary surgery along a torus knot. *Pacific J. Math.* **38** (1971) 737–745.
- [10] K. Murasugi. On the braid index of alternating links. *Trans. Amer. Math. Soc.* **326** (1991) 237–260.

DEPARTMENT OF MATHEMATICS AND STATISTICS, CALIFORNIA STATE UNIVERSITY, CHICO, CHICO
CA 95929-0525, USA

E-mail address: TMattman@CSUChico.edu

E-mail address: GMaybrun@digitalpath.net

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, MATHEMATICS, BLDG. 380, 450 SERRA
MALL STANFORD, CA 94305-2125, USA

E-mail address: krob321@stanford.edu